

THE PAVING CONJECTURE IS EQUIVALENT TO THE PAVING CONJECTURE FOR TRIANGULAR MATRICES

PETER G. CASAZZA AND JANET C. TREMAIN

ABSTRACT. We resolve a 25 year old problem by showing that The Paving Conjecture is equivalent to The Paving Conjecture for Triangular Matrices.

1. INTRODUCTION

The Kadison-Singer Problem [14] has been one of the most intractable problems in mathematics for nearly 50 years.

Kadison-Singer Problem (KS). *Does every pure state on the (abelian) von Neumann algebra $\mathbb{D}$ of bounded diagonal operators on ℓ_2 have a unique extension to a (pure) state on $B(\ell_2)$, the von Neumann algebra of all bounded linear operators on the Hilbert space ℓ_2 ?*

A **state** of a von Neumann algebra $\mathcal{R}$ is a linear functional f on $\mathcal{R}$ for which $f(I) = 1$ and $f(T) \geq 0$ whenever $T \geq 0$ (i.e. whenever T is a positive operator). The set of states of $\mathcal{R}$ is a convex subset of the dual space of $\mathcal{R}$ which is compact in the w^* -topology. By the Krein-Milman theorem, this convex set is the closed convex hull of its extreme points. The extremal elements in the space of states are called the **pure states** (of $\mathcal{R}$). The Kadison-Singer Problem had been dormant for many years when it was recently brought back to life in [9] and [10] where it was shown that KS is equivalent to fundamental unsolved problems in a dozen different areas of research in pure mathematics, applied mathematics and engineering.

A significant advance on the Kadison-Singer Problem was made by Anderson [2] in 1979 when he reformulated KS into what is now known as the **Paving Conjecture** (Lemma 5 of [14] shows a connection between KS and Paving). Before we state this conjecture, let us introduce some notation. For an operator T on ℓ_2^n , its matrix representation $(\langle Te_i, e_j \rangle)_{i,j \in I}$ is with respect to the natural orthonormal basis. If $A \subset \{1, 2, \dots, n\}$, the **diagonal projection** Q_A is the matrix all of whose entries are zero except for the (i, i) entries for $i \in A$ which are all one.

1991 *Mathematics Subject Classification.* Primary: 47A20, 47B99; Secondary: 46B07.
The first author was supported by NSF DMS 0405376.

Paving Conjecture (PC). *For $\epsilon > 0$, there is a natural number r so that for every natural number n and every linear operator T on l_2^n whose matrix has zero diagonal, we can find a partition (i.e. a paving) $\{A_j\}_{j=1}^r$ of $\{1, \dots, n\}$, so that*

$$\|Q_{A_j} T Q_{A_j}\| \leq \epsilon \|T\| \quad \text{for all } j = 1, 2, \dots, r.$$

It is important that r not depend on n in PC. We will say that an arbitrary operator T satisfies PC if $T - D(T)$ satisfies PC where $D(T)$ is the diagonal of T . It is known that the class of operators satisfying PC (the **pavable operators**) is a closed subspace of $B(l_2)$. Also, to verify PC we only need to verify it for any one the following classes of operators [1, 10, 8]: 1. unitary operators, 2. positive operators, 3. orthogonal projections (or just orthogonal projections with $1/2$'s on the diagonal), 4. Gram operators of the form $T^*T = (\langle f_i, f_j \rangle)_{i,j \in I}$ where $\|f_i\| = 1$ and $Te_i = f_i$ is a bounded operator. The only large classes of operators which have been shown to be pavable are “diagonally dominant” matrices [3, 4, 12], matrices with all entries real and positive [5, 13] and matrices with small entries [6].

Since the beginnings of the *paving era*, it has been a natural question whether PC is equivalent to PC for triangular operators. This question was formally asked several times at meetings by Gary Weiss and Lior Tzafriri and appeared (for a short time) on the AIM website (<http://www.aimath.org/The Kadison-Singer Problem>) as an important question for PC. In this paper we will verify this conjecture. Given two conjectures C_1, C_2 we say that C_1 **implies** C_2 if a positive answer to C_1 implies a positive answer for C_2 . They are **equivalent** if they imply each other.

2. PRELIMINARIES

Recall that a family of vectors $\{f_i\}_{i \in I}$ is a **Riesz basic sequence** in a Hilbert space $\mathbb{H}$ if there are constants $A, B > 0$ so that for all scalars $\{a_i\}_{i \in I}$ we have:

$$A^2 \sum_{i \in I} |a_i|^2 \leq \left\| \sum_{i \in I} a_i f_i \right\|^2 \leq B^2 \sum_{i \in I} |a_i|^2.$$

We call A, B the **lower and upper Riesz basis bounds** for $\{f_i\}_{i \in I}$. If $\epsilon > 0$ and $A = 1 - \epsilon, B = 1 + \epsilon$ we call $\{f_i\}_{i \in I}$ an **ϵ -Riesz basic sequence**. If $\|f_i\| = 1$ for all $i \in I$ this is a **unit norm Riesz basic sequence**. A natural question is whether we can improve the Riesz basis bounds for a unit norm Riesz basic sequence by partitioning the sequence into subsets.

R_ϵ -Conjecture. *For every $\epsilon > 0$, every unit norm Riesz basic sequence is a finite union of ϵ -Riesz basic sequences.*

The R_ϵ -Conjecture was posed by Casazza and Vershynin [11] where it was shown that KS implies this conjecture. It is now known that the R_ϵ -Conjecture

is equivalent to KS [9]. We will show that PC for triangular operators implies a positive solution to the R_ϵ -Conjecture. Actually, we need the finite dimensional quantative version of this conjecture.

Finite R_ϵ -Conjecture. *Given $0 < \epsilon, A, B$, there is a natural number $r = r(\epsilon, A, B)$ so that for every $n \in \mathbb{N}$ and every unit norm Riesz basic sequence $\{f_i\}_{i=1}^n$ for ℓ_2^n with Riesz basis bounds $0 < A \leq B$, there is a partition $\{A_j\}_{j=1}^r$ of $\{1, 2, \dots, n\}$ so that for all $j = 1, 2, \dots, r$ the family $\{f_i\}_{i \in A_j}$ is an ϵ -Riesz basic sequence.*

There are standard methods for turning infinite dimensional results into quantative finite dimensional results so we will just outline the proof of their equivalence. We will need a proposition from [7].

Proposition 2.1. *Fix a natural number r and assume for every natural number n we have a partition $\{A_i^n\}_{i=1}^r$ of $\{1, 2, \dots, n\}$. Then there are natural numbers $\{n_1 < n_2 < \dots\}$ so that if $j \in A_i^{n_j}$ for some $i \in \{1, \dots, r\}$, then $j \in A_i^{n_k}$, for all $k \geq j$. Hence, if $A_i = \{j \mid j \in A_i^{n_j}\}$ then*

- (1) $\{A_i\}_{i=1}^r$ is a partition of $\mathbb{N}$.
- (2) If $A_i = \{j_1^i < j_2^i < \dots\}$ then for every natural number k we have $\{j_1^i, j_2^i, \dots, j_k^i\} \subset A_i^{n_{j_k^i}}$.

Theorem 2.2. *The R_ϵ -Conjecture is equivalent to the Finite R_ϵ -Conjecture.*

Proof. Assume the Finite R_ϵ -Conjecture is true. Let $\{f_i\}_{i=1}^\infty$ be a unit norm Riesz basic sequence in $\mathbb{H}$ with bounds $0 < A, B$ and fix $\epsilon > 0$. Then there is a natural number $r \in \mathbb{N}$ so that for all $n \in \mathbb{N}$ there is a partition $\{A_j^n\}_{j=1}^r$ of $\{1, 2, \dots, n\}$ and for every $j = 1, 2, \dots, r$ the family $\{f_i\}_{i \in A_j^n}$ is an ϵ -Riesz basic sequence. Choose a partition $\{A_j\}_{j=1}^r$ of $\mathbb{N}$ satisfying Proposition 2.1. By (2) of this proposition, for each $j = 1, 2, \dots, r$, the first n -elements of $\{f_i\}_{i \in A_j}$ come from one of the A_j^n and hence form an ϵ -Riesz basic sequence. So $\{f_i\}_{i \in A_j}$ is an ϵ -Riesz basic sequence.

Now assume the the Finite R_ϵ -Conjecture fails. Then there is some $0 < \epsilon, A, B$, natural numbers $n_1 < n_2 < \dots$ and unit norm Riesz basic sequences $\{f_i^r\}_{i=1}^{n_r}$ for $\ell_2^{n_r}$ so that whenever $\{A_j\}_{j=1}^r$ is a partition of $\{1, 2, \dots, n_r\}$ one of the sets $\{f_i^r\}_{i \in A_j}$ is not an ϵ -Riesz basic sequence. Considering

$$\{f_i\}_{i=1}^\infty = \{f_i^r\}_{i=1, r=1}^{n_r, \infty} \in \left(\sum_{r=1}^\infty \oplus \ell_2^{n_r} \right)^{1/2},$$

we see that this family of vectors forms a unit norm Riesz basic sequence with bounds $0 < A, B$ but for any natural number r and any partition $\{A_j\}_{j=1}^r$ of $\mathbb{N}$ one of the sets $\{f_i\}_{i \in A_j}$ is not an ϵ -Riesz basic sequence. $\square$

3. THE MAIN THEOREM

Our main theorem is:

Theorem 3.1. *The Paving Conjecture is equivalent to the Paving Conjecture for Triangular matrices.*

Proof. Since a paving of T is also a paving of T^* , we only need to show that The Paving Conjecture for Lower Triangular Operators implies the Finite R_ϵ -Conjecture. Fix $0 < \epsilon, A, B$, fix $n \in \mathbb{N}$ and let $\{f_i\}_{i=1}^n$ be a unit norm Riesz basis for ℓ_2^n with bounds A, B . We choose a natural number $r \in \mathbb{N}$ satisfying:

$$1 - \frac{B^4}{A^4 r} \geq 1 - \frac{\epsilon}{2}.$$

We will do the proof in 5 steps.

Step 1: There is a partition $\{A_j\}_{j=1}^r$ of $\{1, 2, \dots, n\}$ so that for every $j = 1, 2, \dots, r$ and every $i \in A_j$ and every $1 \leq k \neq j \leq r$ we have:

$$\sum_{i \neq \ell \in A_j} |\langle f_i, f_\ell \rangle|^2 \leq \sum_{\ell \in A_k} |\langle f_i, f_\ell \rangle|^2.$$

The argument for this is due to Halpern, Kaftal and Weiss ([13], Proposition 3.1) so we will outline it for our case. Out of all ways of partitioning $\{1, 2, \dots, n\}$ into r -sets, choose one, say $\{A_j\}_{j=1}^r$, which minimizes

$$(3.1) \quad \sum_{j=1}^r \sum_{i \in A_j} \sum_{i \neq \ell \in A_j} |\langle f_i, f_\ell \rangle|^2.$$

We now observe that for each $1 \leq j \leq r$, each $i \in A_j$ and all $1 \leq k \neq j \leq r$ we have

$$\sum_{i \neq \ell \in A_j} |\langle f_i, f_\ell \rangle|^2 \leq \sum_{\ell \in A_k} |\langle f_i, f_\ell \rangle|^2.$$

To see this, assume this inequality fails. That is, for some j_0, i_0, k_0 as above we have

$$\sum_{i_0 \neq \ell \in A_{j_0}} |\langle f_{i_0}, f_\ell \rangle|^2 > \sum_{\ell \in A_{k_0}} |\langle f_{i_0}, f_\ell \rangle|^2.$$

We define a new partition $\{B_j\}_{j=1}^r$ of $\{1, 2, \dots, n\}$ by: $B_j = A_j$ if $j \neq j_0, k_0$; $B_{j_0} = A_{j_0} - \{i_0\}$; $B_{k_0} = A_{k_0} \cup \{i_0\}$. It easily follows that

$$\sum_{j=1}^r \sum_{i \in B_j} \sum_{i \neq \ell \in B_j} |\langle f_i, f_\ell \rangle|^2 < \sum_{j=1}^r \sum_{i \in A_j} \sum_{i \neq \ell \in A_j} |\langle f_i, f_\ell \rangle|^2,$$

which contradicts the minimality of Equation 3.1.

Step 2: For every $j = 1, 2, \dots, r$ and every $i \in A_j$ we have

$$\sum_{i \neq \ell \in A_j} |\langle f_i, f_\ell \rangle|^2 \leq \frac{B^2}{r}.$$

Define an operator $Sf = \sum_{i=1}^n \langle f, f_i \rangle f_i$. Then,

$$\langle Sf, f \rangle = \sum_{i=1}^r |\langle f, f_i \rangle|^2,$$

and since $\{f_i\}_{i=1}^n$ is a Riesz basis with bounds A, B we have

$$A^2 I \leq S \leq B^2 I.$$

Now, by Step 1,

$$\begin{aligned} \sum_{i \neq \ell \in A_j} |\langle f_i, f_\ell \rangle|^2 &\leq \frac{1}{r} \left[\sum_{i \neq \ell \in A_j} |\langle f_i, f_\ell \rangle|^2 + \sum_{j \neq k=1}^r \sum_{\ell \in A_k} |\langle f_i, f_\ell \rangle|^2 \right] \\ &\leq \frac{1}{r} \sum_{i=1}^n |\langle f_i, f_\ell \rangle|^2 \\ &\leq \frac{1}{r} \|S\| \|f_i\|^2 \leq \frac{B^2}{r}. \end{aligned}$$

Step 3: For each $j = 1, 2, \dots, r$ and all $i \in A_j$, if P_{ij} is the orthogonal projection of $\text{span } \{f_\ell\}_{\ell \in A_j}$ onto $\text{span } \{f_\ell\}_{i \neq \ell \in A_j}$ then

$$\|P_{ij} f_i\|^2 \leq \frac{B^4}{A^4 r}.$$

Define the operator S_{ij} on $\text{span } \{f_\ell\}_{\ell \in A_j}$ by

$$S_{ij}(f) = \sum_{i \neq \ell \in A_j} \langle f, f_\ell \rangle f_\ell.$$

Then $A^2 I \leq S_{ij} \leq B^2 I$ and $\{S_{ij}^{-1} f_\ell\}_{i \neq \ell \in A_j}$ are the dual functionals for the Riesz basic sequence $\{f_\ell\}_{i \neq \ell \in A_j}$. Also, as in Step 1, $A^2 I \leq S_{ij} \leq B^2 I$. So by

Step 2,

$$\begin{aligned}
\|P_{ij}f_i\|^2 &= \left\| \sum_{i \neq \ell \in A_j} \langle f_i, f_\ell \rangle S_{ij}^{-1} f_\ell \right\|^2 \\
&\leq \|S_{ij}^{-1}\|^2 \left\| \sum_{i \neq \ell \in A_j} \langle f_i, f_\ell \rangle f_\ell \right\|^2 \\
&\leq \frac{B^2}{A^4} \sum_{i \neq \ell \in A_j} |\langle f_i, f_\ell \rangle|^2 \leq \frac{B^4}{A^4 r}.
\end{aligned}$$

Step 4: Fix $1 \leq j \leq r$ and let $A_j = \{i_1, i_2, \dots, i_k\}$. If we Gram-Schmidt $\{f_{i_\ell}\}_{\ell=1}^k$ to produce an orthonormal basis $\{e_{i_\ell}\}_{\ell=1}^k$ then for all $1 \leq m \leq k$ we have

$$|\langle f_{i_m}, e_{i_m} \rangle|^2 \geq 1 - \frac{\epsilon}{2}.$$

Fix $1 \leq m \leq k$ and let Q_m be the orthogonal projection of $\text{span } \{e_{i_\ell}\}_{\ell=1}^k$ onto $\text{span } \{e_{i_\ell}\}_{\ell=1}^m = \text{span } \{f_{i_\ell}\}_{\ell=1}^m$. By Step 3,

$$\|Q_m f_{i_m}\|^2 \leq \|P_{mj} f_{i_m}\|^2 \leq \frac{B^4}{A^4 r}.$$

Since

$$f_{i_m} = \sum_{\ell=1}^m \langle f_{i_\ell}, e_{i_\ell} \rangle e_{i_\ell},$$

we have

$$\begin{aligned}
|\langle f_{i_m}, e_{i_m} \rangle|^2 &= \|f_{i_m}\|^2 - \|Q_{m-1} f_{i_m}\|^2 \\
&\geq 1 - \frac{B^4}{A^4 r} \geq 1 - \frac{\epsilon}{2},
\end{aligned}$$

where the last inequality follows from our choice of r .

Step 5: We complete the proof.

Let

$$M = (\langle f_{i_s}, e_{i_t} \rangle)_{s \neq t=1}^k,$$

where by this notation we mean the $k \times k$ -matrix with zero diagonal and the given values off the diagonal. By the Gram-Schmidt Process, M is a lower triangular matrix with zero diagonal. Define an operator $T : \ell_2^k \rightarrow \text{span } \{e_{i_\ell}\}_{\ell=1}^k$ by

$$T((a_{i_\ell})_{\ell=1}^k) = \sum_{\ell=1}^k a_{i_\ell} f_{i_\ell}.$$

If K is the matrix of T with respect to the orthonormal basis $\{e_{i_\ell}\}_{\ell=1}^k$ and $D = D(K)$ is the diagonal of K then $M = (K - D)^*$ and so

$$\|M\| \leq \|K\| + \|D\| = \|T\| + 1 \leq B + 1.$$

By The Paving Conjecture for lower triangular matrices, there is a natural number L_j (which is a function of $0 < \epsilon$ and B only) and a partition $\{B_\ell^j\}_{\ell=1}^{L_j}$ of $\{i_1, i_2, \dots, i_k\}$ so that

$$\|Q_{B_\ell^j} M Q_{B_\ell^j}\| \leq \frac{\epsilon}{2},$$

for all $\ell = 1, 2, \dots, L_j$ ($Q_{B_\ell^j}$ was defined in the introduction). Now, for all scalars $(a_{i_s})_{i_s \in B_\ell^j}$, if

$$f = \sum_{i_s \in B_\ell^j} a_{i_s} f_{i_s},$$

then

$$\begin{aligned} \left\| \sum_{i_s \in B_\ell^j} a_{i_s} f_{i_s} \right\| &= \|D(f) + Q_{B_\ell^j} M^* Q_{B_\ell^j}(f)\| \\ &\geq \|Df\| - \|Q_{B_\ell^j} M^* Q_{B_\ell^j}(f)\| \\ &\geq (1 - \frac{\epsilon}{2})\|f\| - \frac{\epsilon}{2}\|f\| \\ &\geq (1 - \epsilon)\|f\|. \end{aligned}$$

Similarly,

$$\left\| \sum_{i_s \in B_\ell^j} a_{i_s} f_{i_s} \right\| \leq (1 + \epsilon)\|f\|.$$

It follows that $\{f_i\}_{i \in B_\ell^j}$ is an ϵ -Riesz basic sequence for all $j = 1, 2, \dots, r$ and all $\ell = 1, 2, \dots, L_j$. Hence, the Finite R_ϵ -Conjecture holds which completes the proof of the theorem. $\square$

Let us make an observation concerning the proof of the main theorem.

Definition 3.2. Let $\{f_i\}_{i=1}^\infty$ be a sequence of vectors in a Hilbert space $\mathbb{H}$. For each $i = 1, 2, \dots$ let P_i be the orthogonal projection of $\mathbb{H}$ onto $\text{span}\{f_\ell\}_{\ell \neq i \in \mathbb{N}}$. Our sequence is said to be ϵ -**minimal** if $\|P_i\| \leq \epsilon$ for all $i = 1, 2, \dots$.

The first three steps of the proof of Theorem 3.1 yields:

Corollary 3.3. If $\{f_i\}_{i=1}^\infty$ is a unit norm Riesz basic sequence in a Hilbert space $\mathbb{H}$ then for every $\epsilon > 0$ there is a partition $\{A_j\}_{j=1}^r$ of $\mathbb{N}$ so that for all $j = 1, 2, \dots, r$, the family $\{f_i\}_{i \in A_j}$ is ϵ -minimal.

REFERENCES

- [1] C.A. Akemann and J. Anderson, *Lyapunov theorems for operator algebras*, Mem. AMS **94** (1991).
- [2] J. Anderson, *Extreme points in sets of positive linear maps on $B(\mathbb{H})$* , Jour. Functional Analysis **31** (1979) 195–217.
- [3] R. Balan, P.G. Casazza, C. Heil and Z. Landau, *Density, overcompleteness and localization of frames. I. Theory*, Preprint.
- [4] R. Balan, P.G. Casazza, C. Heil and Z. Landau, *Density, overcompleteness and localization of frames. II. Gabor systems*, Preprint.
- [5] K. Berman, H. Halpern, V. Kaftal and G. Weiss, *Matrix norm inequalities and the relative Dixmier property*, Integ. Eqns. and Operator Theory **11** (1988) 28–48.
- [6] J. Bourgain and L. Tzafriri, *Invertibility of “large” submatrices and applications to the geometry of Banach spaces and Harmonic Analysis*, Israel J. Math. **57** (1987) 137–224.
- [7] P.G. Casazza, O. Christensen, A. Lindner and R. Vershynin, *Frames and the Feichtinger conjecture*, Proceedings of AMS, **133** No. 4 (2005) 1025–1033.
- [8] P.G. Casazza, D. Edidin, D. Kalra and V. Paulsen, *The Kadison-Singer Problem and Projections*, Preprint.
- [9] P.G. Casazza and J.C. Tremain, *The Kadison-Singer Problem in Mathematics and Engineering*, Proceedings of the National Academy of Sciences, **103** No. 7 (2006) 2032–2039.
- [10] P.G. Casazza, M. Fickus, J.C. Tremain, and E. Weber, *The Kadison-Singer Problem in Mathematics and Engineering: Part II: A detailed account*. (Accepted for The Proceedings of The 2005 Great Plains Operator Theory Symposium (GPOTS), Contemp. Math., Amer. Math. Soc., to appear in 2006).
- [11] P.G. Casazza and R. Vershynin, *Kadison-Singer meets Bourgain-Tzafriri*, Preprint.
- [12] K.H. Gröchenig, *Localized frames are finite unions of Riesz sequences*, Adv. Comp. Math. **18** (2003) 149–157.
- [13] H. Halpern, V. Kaftal and G. Weiss, *Matrix pavings and Laurent operators*, J. Op. Th. **16** (1986) 121–140.
- [14] R. Kadison and I. Singer, *Extensions of pure states*, American Jour. Math. **81** (1959), 383–400.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211-4100

E-mail address: janet,pete@math.missouri.edu